

Frederic A. Johnson

A NATURAL DEDUCTION RELEVANCE LOGIC

The relevance logic (NDR) presented in this paper is the result of an attempt to find a natural deduction development, in the style of I. M. Copi (**Introduction to Logic**, 4th ed., MacMillan, 1972), for the relevance logic I presented in “A Three-Valued Interpretation for a Relevance Logic” (**The Relevance Logic Newsletter**, Vol. 1, no. 3, 1976).

The propositional variables of NDR are, p_1, p_2, \dots . NDR’s well-formed formulas are constructed in the standard way by using propositional variables, parentheses and the connectives, $-$, \cdot and \vee , in order of increasing binding strength. ‘ $P \supset Q$ ’ is by definition ‘ $-(P \cdot -Q)$ ’. Capital letters with or without subscripts are metalinguistic variables which range over the well-formed formulas. We will use ‘ \vdash_r ’ to present NDR’s rules of inference:

1. $P \vdash_r P \vee Q$, where every p_i in Q occurs in P . (Restricted Addition, RA)
2. $P \vdash_r P \cdot (Q \vee -Q)$, where every p_i in Q occurs in P . (Restricted Tautology Conjunction, RTC)
3. $P, Q \vdash_r P \cdot Q$ (Conjunction, Conj.)
4. $P \cdot Q \vdash_r P$ (Simplification, Simp.)
5. $P \vee Q \cdot R \vdash_r P \vee Q$ (Disjunctive Simplification, DS)
6. $P \vee Q \cdot -Q \vdash_r P$ (Contradiction Elimination, CE)
7. If $S \equiv_l T$ in virtue of exactly one of the following statements then $F(S) \vdash F(T)$.
 - i) $P \cdot (Q \vee R) \equiv_l P \cdot Q \vee P \cdot R$ (DeMorgan’s, DeM)
 - ii) $P \cdot (Q \vee R) \equiv_l P \cdot Q \vee P \cdot R$ (Distribution, Dist.)
 $P \vee Q \cdot R \equiv_l (P \vee Q) \cdot (P \vee R)$

- | | | |
|------|--|-----------------------|
| iii) | $P \cdot (Q \cdot R) \equiv_l (P \cdot Q) \cdot R$
$P \vee (Q \vee R) \equiv_l (P \vee Q) \vee R$ | (Association, Assoc.) |
| iv) | $P \cdot Q \equiv_l Q \cdot P$
$P \vee Q \equiv_l Q \vee P$ | (Computation, Com.) |
| v) | $--P \equiv_l P$ | (Double Negation, DN) |
| vi) | $P \cdot P \equiv_l P$
$P \vee P \equiv_l P$ | (Tautology, Taut.) |

NDR's entailment relation, symbolized by ' \vdash ', is defined as follows: $P_1, \dots, P_n \vdash C$ if and only if there is a sequence of well-formed formulas S_1, \dots, S_m such that $S_m = C$ and each S_i ($1 \leq i \leq m$) is either a P_i ($1 \leq i \leq n$) or follows from preceding S_j by one of the rules of inference.

THEOREM 1. *If $P_1, \dots, P_n \vdash C$ then P_1, \dots, P_n classically entails C and every p_i in C occurs in P_1, \dots, P_n .*

PROOF. Every valuation which assigns t to the premises of the rules of inference assigns t to the conclusion. Furthermore, none of the rules of inference introduce into the conclusion propositional variables which do not occur in the premises.

THEOREM 2. (Indirect Proof.) *If $P \cdot -Q \vdash R \cdot -R$ and every p_i in Q occurs in P then $P \vdash Q$.*

PROOF. Let S_1, \dots, S_n be a sequence of well-formed formulae such that $S_1 = P \cdot -Q$, $S_n = R \cdot -R$ and each S_i ($1 \leq i \leq n$) is either $P \cdot -Q$ or follows from S_j or from S_j and S_k ($1 \leq j, k < n$). Then construct this sequence of statements:

- | | | |
|--------------|-----------------------------|---------------------------------|
| 1. | P | |
| 2. | $P \cdot (Q \vee -Q)$ | 1, RTC |
| $a_1(= 3)$. | $P \cdot Q \vee P \cdot -Q$ | $(P \cdot S \vee S_1)$ 2, Dist. |
| | ⋮ | |
| | ⋮ | |
| | ⋮ | |
| a_2 . | $P \cdot Q \vee S_2$ | |
| | ⋮ | |
| | ⋮ | |
| | ⋮ | |
| a_n . | $P \cdot Q \vee S_n$ | |

$a_n + 1.$	$P \cdot Q$	a_n , CE
$a_n + 2.$	$Q \cdot P$	$a_n + 1$, Com.
$a_n + 3.$	Q	$a_n + 2$, Simp.

The steps from, but excluding, $P \cdot Q \vee S_{j-1}$ to, and including, $P \cdot Q \vee S_j$ for $1 < j \leq n$ are to be filled in as follows:

- i) If $S_j = P \cdot \neg Q$ then supply the sequence
- | | | |
|------------|---|-------------------|
| $a_j - 1.$ | $(P \cdot Q \vee P \cdot \neg Q) \cdot (Q \vee \neg Q)$ | a_1 , RTC |
| $a_j.$ | $P \cdot Q \vee P \cdot \neg Q$ | $a_j - 1$, Simp. |
- Make $a_j - 2 = a_{j-1}$.
- ii) If $S_i \vdash S_j$ ($i < j$) by RA, where $S_j = S_i \vee T$, then supply the sequence
- | | | |
|------------|-------------------------------|--------------------|
| $a_j - 1.$ | $(P \cdot Q \vee S_i) \vee T$ | a_i , RA |
| $a_j.$ | $P \cdot Q \vee (S_i \vee T)$ | $a_j - 1$, Assoc. |
- Make $a_j - 2 = a_{j-1}$.
- iii) If $S_i \vdash S_j$ ($i < j$) by RTC, where $S_j = S_i \cdot (T \vee \neg T)$, then supply the sequence
- | | | |
|------------|---|-------------------|
| $a_j - 7.$ | $(P \cdot Q \vee S_i) \cdot (T \vee \neg T)$ | a_i , RTC |
| $a_j - 6.$ | $(T \vee \neg T) \cdot (P \cdot Q \vee S_i)$ | $a_j - 7$, Com. |
| $a_j - 5.$ | $(T \vee \neg T) \cdot (P \cdot Q) \vee$
$(T \vee \neg T) \cdot S_i$ | $a_j - 6$, Dist. |
| $a_j - 4.$ | $(T \vee \neg T) \cdot S_i \vee (T \vee \neg T) \cdot$
$(P \cdot Q)$ | $a_j - 5$, Com. |
| $a_j - 3.$ | $(T \vee \neg T) \cdot S_i \vee (P \cdot Q) \cdot$
$(T \vee \neg T)$ | $a_j - 4$, Com. |
| $a_j - 2.$ | $(T \vee \neg T) \cdot S_i \vee (P \cdot Q)$ | $a_j - 3$, DS |
| $a_j - 1.$ | $(P \cdot Q) \vee (T \vee \neg T) \cdot S_i$ | $a_j - 2$, Com. |
| $a_j.$ | $(P \cdot Q) \vee S_i \cdot (T \vee \neg T)$ | $a_j - 1$, Com. |
- Make $a_j - 8 = a_{j-1}$.
- iv) If $S_h, S_i \vdash S_j$ ($h, i < j$) by Conj., where $S_j = S_h \cdot S_i$, then supply the sequence
- | | | |
|------------|---|-------------------|
| $a_j - 1.$ | $(P \cdot Q \vee S_h) \cdot (P \cdot Q \vee S_i)$ | a_h, a_i Conj. |
| $a_j.$ | $P \cdot Q \vee (S_h \cdot S_i)$ | $a_j - 1$, Dist. |
- Make $a_j - 2 = a_{j-1}$.

Procedures for filling in the lines between a_j and a_{j-1} when $S_i \vdash S_j$ in virtue of Rules 4-7 are also easily constructed.

THEOREM 3. (Transitivity of Entailment.) *If $P \vdash Q$ and $Q \vdash R$ then $P \vdash R$.*

PROOF. Let $S_1 (= P), S_2, \dots, S_m (= Q)$ be a sequence of well-formed formulas which shows that $P \vdash Q$ and let $S_m (= Q), S_{m+1}, \dots, S_n (= R)$ be a sequence of well-formed formulas which shows that $P \vdash R$. Then S_1, \dots, S_n shows that $P \vdash R$.

THEOREM 4. *If P classically entails Q and every p_i in Q occurs in P then $P \vdash Q$.*

PROOF. Assume the antecedent. Then $P \cdot \neg Q$ is a contradiction. By DeM, Dist., Assoc., Com., DN and Taut. $P \cdot \neg Q \vdash R_1 \cdot \neg R_1 \cdot S_1 \vee \dots \vee R_n \cdot \neg R_n \cdot S_n \cdot (R_1 \cdot \neg R_1 \cdot S_1 \vee \dots \vee R_n \cdot \neg R_n \cdot S_n)$ is one of the formulas which will be produced when following some of the various mechanical procedures for generating the disjunctive normal form of $P \cdot \neg Q$. By CE and Simp. $R_1 \cdot \neg R_1 \cdot S_1 \vee \dots \vee R_n \cdot \neg R_n \cdot S_n \vdash R_1 \cdot \neg R_1$. By Theorem 3 (Th. 3), $P \cdot \neg Q \vdash R_1 \cdot \neg R_1$. By Th. 2 $P \vdash Q$.

THEOREM 5. (Adjunction). *If $P \vdash Q$ and $P \vdash R$ then $P \vdash Q \cdot R$.*

PROOF. Let $S_1, \dots, S_m (= Q), \dots, S_n (= R)$, where $m \leq n$, be a sequence that shows that $P \vdash Q$ and $P \vdash R$. Let $S_{n+1} = Q \cdot R$. Then S_1, \dots, S_{n+1} shows that $P \vdash Q \cdot R$, using Conj.

THEOREM 6. (Deduction Theorem). *If $P \cdot Q$ and every p_i in Q occurs in P then $P \vdash Q \supset C$.*

PROOF. Assume the antecedent. By Theorem 1 $P \cdot Q$ classically entails C . Then P classically entails $Q \supset C$. Since every p_i in Q occurs in P and every p_i in C occurs in $P \cdot Q$ it follows that every p_i in $Q \supset C$ occurs in P . By Theorem 4 $P \vdash Q \supset C$.¹

THEOREM 7. (Antilogism). *If $P \cdot Q \vdash R$ and every p_i in Q occurs in P then $P \cdot \neg R \vdash \neg Q$.*

PROOF. By Simp. $P \cdot \neg R \vdash P$. Assume the antecedent. By Th. 6 and the definition of ' \supset ' $P \vdash \neg(Q \cdot \neg R)$. By Th. 3 $P \cdot \neg R \vdash \neg(Q \cdot \neg R)$. By Com.

¹This proof, suggested by Richard Routley, is more straightforward than my original proof. I am grateful for Professor Routley's comments, which led to several improvements.

and Simp. $P \cdot -R \vdash -R$. By Th. 5 $P \cdot -R \vdash -R \cdot -(Q \cdot -R)$. By Dem, Dist., Com. and Simp. $-R \cdot -(Q \cdot -R) \vdash -Q$. By Th. 3 $P \cdot -R \vdash -Q$.

The difference between NDR and the relevance logic presented in “A Three-Valued Interpretation of a Relevance Logic” is that the latter does not recognize the validity of any arguments with contradictory premises, whereas NDR does. For example, $p_1 \cdot -p_1 \vdash p_1$ in NDR. But both of these logics endorse what W. T. Parry (The Logic of C. I. Lewis’, **The Philosophy of C. I. Lewis**, ed. P. A. Schilpp, 1968, pp. 115–54) called the Proscriptive Principle, which keeps those arguments which contain a p_i that occurs in the conclusion but not in a premise from being valid. Charles Kielkopf (‘Adjunction and Paradoxical Derivations’, **Analysis**, Vol. 35, no. 4, 1975, pp. 127–9) showed that the system which Parry based on the Proscriptive Principle inadvertently permits the derivation of any statement from a contradiction.

Perhaps the most worrisome feature of NDR is that it denies that in general if A entails B then $-B$ entails $-A$. For example, though $p_1 \cdot p_2$ entails p_1 it is false that $-p_1$ entails $-(p_1 \cdot p_2)$. But the reservations which beginning students of logic have about the validity of Unrestricted Addition, which would guarantee that $-p_1$ entails $-p_1 \vee -p_2$ suggest that this apparent defect may be a virtue.²

*Department of Philosophy
Colorado State University
Fort Collins, Colorado 80523*

²I am grateful to Professor Charles Kielkopf and Professor Patrick McKee for their helpful comments.